

Coherent State and Phase Space Descriptions

Jhih-Sheng Wu

2023

Contents

1	Phase Space Pictures	2
2	Coherent States	4
2.1	Quantum Phase	4
2.2	Coherent States	8
2.3	Displaced Vacuum States	10
2.4	Dynamics of Coherent States	12
2.5	Properties of Coherent States	13
2.5.1	Orthogonality	13
2.5.2	Identity	13
2.5.3	Coherent State Representations of Operators	14
3	Phase Space Distributions	15
3.1	Wigner Distribution	15
3.2	Glauber–Sudarshan P -function	16
3.3	Q -function	17

1 Phase Space Pictures

The state of a classical particle is fully determined by its x and p . A useful way to represent the states is the phase space (x, p) , where the horizontal axis is x and the vertical axis is p . A state of a classical particle is one point in the phase space. The time evolution of a state is the trajectory in the phase space. The trajectory $(x(t), p(t))$ contains all the information of the particle. The classic example is the harmonic oscillator with

$$x(t) = x_0 \cos(\omega t + \phi), \quad (1.1)$$

$$p(t) = -\omega x_0 \sin(\omega t + \phi), \quad (1.2)$$

or in the dimensionless expression

$$\tilde{x}(t) = \frac{x(t)}{x_0} = \cos(\omega t + \phi), \quad (1.3)$$

$$\tilde{p}(t) = \frac{p(t)}{\omega x_0} = -\sin(\omega t + \phi). \quad (1.4)$$

The state travels along the trajectory is a unit circle (see Fig. 1).




Figure 1: A classical state is a point in the phase space. The motion of a state is a trajectory. In the case of a harmonic oscillator, the trajectory is a circle.

An ensemble of classical particles are described by the phase space probability density function $f(x, p)$, where to find a particle with a position x and a momentum p is given by

$$f(x, p) dx dp, \quad (1.5)$$

and the normalization condition is

$$\int dx \int dp f(x, p) dx dp = 1. \quad (1.6)$$

Classically, the function $f(x, p)$ of a pure state, i.e., a single particle, is a delta function $f(x, p) = \delta(x - x_0)\delta(p - p_0)$. We have make the analogies $x \leftrightarrow X$ and

$y \leftrightarrow Y$. One question arises: can we define a function similar to $f(x, p)$ to describe states or ensembles of photons? The problem is that a quantum state can not have exact X and Y at the same time. Thus, a quantum state is not a single point in the phase space. Recall the relations

$$X = \frac{a + a^\dagger}{2}, \quad (1.7)$$

$$Y = \frac{a - a^\dagger}{2i}. \quad (1.8)$$

For a coherent state $|\alpha\rangle$, we have the relations

$$\langle X \rangle = \frac{\alpha + \alpha^*}{2}, \quad (1.9)$$

$$\langle Y \rangle = \frac{\alpha - \alpha^*}{2i}. \quad (1.10)$$

As you can show $\sigma(X) = \sigma(Y) = 1/2$ for a coherent state, it means that a state in the phase space is not a point but a blurred circular cloud (see Fig. 2). The size of the cloud reflects the uncertainty relations. Coherent states are the states satisfying the minimum uncertainty relations. In general, an arbitrary state can have a very broad distribution in the phase space.




Figure 2: A coherent state is a fuzzy circle in the phase space.

A mapping of a state $|\psi\rangle$ or an ensemble to a distribution in the phase space (X, Y) (or equivalently the complex α space.) provide a physical picture. However, a mapping $|\psi\rangle \rightarrow f(X, Y)$ is not uniquely defined. The problem comes from that X and Y are non-commutative operators. There exist many attempts to define a probability density $f(X, Y)$ or $f(\alpha)$. We are going to introduce the three most used definitions,

- Wigner distribution
- Q-function
- P-function

Note that the definitions and calculations of these functions are quite mathematically involved. These functions serve as quantitative tools to describe the phase space probability densities. It is fine to have a qualitative picture in mind first and know more calculations when it is needed.

2 Coherent States

We have shown that the number states $|n\rangle$ do not behave similarly as the classical fields. For example, the expectation value $\langle n|\hat{E}|n\rangle$ is not only static but also zero. A classical field is a field whose amplitude is a harmonic function of t , i.e., $\exp(\pm i\omega t)$. Since the number states form a complete set of the basis vectors, all the photon states, including the classical field, can be written in the number state basis. Hence, we write a classical field $\mathbf{E}_{\text{cl}}(\mathbf{r}, t)$ as a superposition of the number states,

$$|\text{classical}\rangle = \sum_n C_n |n(t)\rangle = \sum_n C_n e^{-in\omega t} |n(0)\rangle. \quad (2.1)$$

The coefficients C_n are to be determined to satisfy the following properties. The classical field $\mathbf{E}_{\text{cl}}(\mathbf{r}, t)$ is the expectation value of the electric field of the classical state,

$$\mathbf{E}_{\text{cl}}(\mathbf{r}, t) = \langle \text{classical} | \hat{\mathbf{E}} | \text{classical} \rangle, \quad (2.2)$$

where for a mode of frequency ω , the classical field $\mathbf{E}_{\text{cl}}(\mathbf{r}, t)$ is sinusoidal,

$$\mathbf{E}_{\text{cl}}(\mathbf{r}, t) = \mathcal{E}_\omega(\mathbf{r}) e^{-i\omega t + \phi}. \quad (2.3)$$

A classical field has the two features, the harmonic oscillation term $e^{-i\omega t}$ and the phase ϕ . Although the expectation value by Eq. (2.3) define the exact values of the amplitude and the phase, the amplitude and phase of the electric field of a state $|\psi\rangle$ in general have uncertainties. Hence, the amplitude and phase of a state should be described by probability distributions.

Note 1: Coherent State

A coherent state is a most classical state of which the amplitude is a finite constant, the phase grows as ωt , and the uncertainties of the amplitude and phase are minimized.

Below, we first discuss how to obtain the phase distribution of a state $|\psi\rangle$, and find the coefficient C_n of a coherent state.

2.1 Quantum Phase

In quantum optics, the electric field \mathbf{E} of an arbitrary photon state $|\psi\rangle$ has the uncertainties in both its amplitude and phase, that is, $\langle E^2 \rangle \neq 0$ and $\langle \phi^2 \rangle \neq 0$.

Indeed, we have not talked about how to obtain ϕ of a photon state $|\psi\rangle$. Note that the phase ϕ is not the phase of a wavefunction but the phase of the electric field. Since \mathbf{E} is an operator but not a number, it turns out that there are many definitions of the phase ϕ . Moreover, the phase ϕ of a state $|\psi\rangle$ is not a single value but a distribution with a finite variance. We will define a phase distribution $\mathcal{P}(\phi)$ where $\mathcal{P}(\phi)d\phi$ is the probability to find the state to have a phase ϕ . Here, we follow the approach by Susskind and Glogower to obtain the phase distribution. The Susskind–Glogower operators are defined by

$$A \equiv (aa^\dagger)^{-\frac{1}{2}}a = (N+1)^{-\frac{1}{2}}a, \quad (2.4)$$

$$A^\dagger \equiv a^\dagger(aa^\dagger)^{-\frac{1}{2}} = a^\dagger(N+1)^{-\frac{1}{2}}. \quad (2.5)$$

If we temporarily treat a as a complex number, $a = |a|\exp i\phi$, the operator A will look as $A = \exp i\phi$. This is the motivation of the definitions, which is to make the operator A taking out the phase factor $\exp i\phi$ of a state. The properties of the SG operators are

$$A|n\rangle = \begin{cases} |n-1\rangle, & n \neq 0, \\ 0, & n = 0, \end{cases} \quad (2.6)$$

$$A^\dagger|n\rangle = |n+1\rangle, \quad (2.7)$$

in the number state bases,

$$A = \sum_n |n\rangle\langle n+1|, \quad (2.8)$$

$$A^\dagger = \sum_n |n+1\rangle\langle n|, \quad (2.9)$$

$$AA^\dagger = 1, \quad (2.10)$$

$$A^\dagger A = 1 - |0\rangle\langle 0|. \quad (2.11)$$

The eigenstate of A is $|\phi\rangle$,

$$A|\phi\rangle = e^{i\phi}|\phi\rangle. \quad (2.12)$$

The state $|\phi\rangle$ in the number states is

$$|\phi\rangle = \sum_n e^{in\phi}|n\rangle. \quad (2.13)$$

The state given by Eq. (2.13) is not normalized. The states $|\phi\rangle$ and $|\phi'\rangle$ are not orthogonal, that is, $\langle\phi'|\phi\rangle \neq 0$. Using the fact

$$\frac{1}{2\pi} \int_0^{2\pi} e^{i(n-n')\phi} d\phi = \delta_{n,n'}, \quad (2.14)$$

we can show that

$$\frac{1}{2\pi} \int_0^{2\pi} d\phi |\phi\rangle\langle\phi| = \mathbb{1}. \quad (2.15)$$

Derivation 1: Identity with Phase States

Let $|\psi\rangle$ be an arbitrary state. In the number state bases, it is

$$|\psi\rangle = \sum_n C_n |n\rangle. \quad (2.16)$$

Applying the operator in Eq. (2.15) on the state, we obtain

$$\frac{1}{2\pi} \int_0^{2\pi} d\phi |\phi\rangle \langle \phi | \psi\rangle = \sum_{C_n} \frac{1}{2\pi} \int_0^{2\pi} d\phi |\phi\rangle \langle \phi | n\rangle \quad (2.17)$$

$$= \frac{1}{2\pi} \sum_n \int d\phi |\phi\rangle C_n e^{-in\phi} \quad (2.18)$$

$$= \frac{1}{2\pi} \sum_{n,m} \int d\phi e^{im\phi} |m\rangle C_n e^{-in\phi} \quad (2.19)$$

$$= \sum_{n,m} \delta_{mn} C_n |m\rangle \quad (2.20)$$

$$= \sum_n C_n |n\rangle \quad (2.21)$$

$$= |\psi\rangle, \quad (2.22)$$

which proves the operator in Eq. (2.15) is an identity.

The phase distribution $\mathcal{P}(\phi)$ of a state $|\psi\rangle$ is

$$\mathcal{P}(\phi) \equiv \frac{1}{2\pi} |\langle \phi | \psi\rangle|^2 \quad (2.23)$$

$$= \frac{1}{2\pi} \left| \sum_n C_n e^{-in\phi} \right|^2. \quad (2.24)$$

The phase distribution $\mathcal{P}(\phi)$ is normalized,

$$\int_0^{2\pi} \mathcal{P}(\phi) d\phi = 1. \quad (2.25)$$

The phase distribution $\mathcal{P}(\phi)$ of an ensemble is

$$\mathcal{P}(\phi) = \frac{1}{2\pi} \langle \phi | \rho | \phi \rangle. \quad (2.26)$$

Note 2: Phase of a Phase State

The phase distribution function $\mathcal{P}(\phi)$ reveals the phase distribution of a state $|\psi\rangle$. Since N and A does not commute ($[N, A] = -A$), a state can not have a single phase but a phase distribution. The phase state $|\phi'\rangle$ is supposed to have a specific phase ϕ' . However, since the phase state is not normalized, it is not physical but a mathematical tool. We consider an approximate phase

state which is normalized,

$$|\phi'\rangle_{\text{app}} \equiv \sum_{n=0}^{N_{\text{max}}} \frac{e^{in\phi'} |n\rangle}{\sqrt{N_{\text{max}} + 1}}. \quad (2.27)$$

The phase distribution function of $|\phi'\rangle_{\text{app}}$ is

$$\mathcal{P}(\phi) = \frac{1}{2(N_{\text{max}} + 1)\pi} \left| \frac{\sin\left[\frac{(N_{\text{max}}+1)(\phi-\phi')}{2}\right]}{\sin\left[\frac{\phi-\phi'}{2}\right]} \right|^2. \quad (2.28)$$




Figure 3: Phase distribution functions of $|\phi_1 = 0.5\pi\rangle$ and $|\phi_2 = 0.7\pi\rangle$. The maximum number is $N_{\text{max}} = 12$.

```

1 import matplotlib
2 import matplotlib.pyplot as plt
3 import numpy as np
4 # Data for plotting
5 phi1 = 0.5 * np.pi
6 phi2 = 0.7 * np.pi
7 Nmax = 12
8 phi = np.arange(0.0, 2.0 * np.pi, 0.01)
9 # define the phase distribution function
10 def phase_dist_func(x,y):
11     return np.sin(Nmax*(x-y)/2)**2./np.sin((x-y)/2)**2/Nmax/(2*np.pi)
12 phase_dist_1 = phase_dist_func(phi,phi1)
13 phase_dist_2 = phase_dist_func(phi,phi2)
14 ## plot
15 fig, ax = plt.subplots()
16 ax.plot(phi, phase_dist_1, label=r'$\phi_1=0.5\pi$')
17 ## r: raw string
18 ax.plot(phi, phase_dist_2, label=r'$\phi_2=0.7\pi$')
19 ## r: raw string
20 ax.set(xlabel='$\phi$', ylabel='$\mathcal{P}(\phi)$',
21       title='Phase Distribution Function of a Phase State')
22 ax.grid()
23 plt.legend()
24 fig.savefig("phase_dist.png", dpi=300)
25 plt.show()

```

Figure 4: Python codes.

Exercise 1: Phase Distribution Function

Show Eq. (2.28). Use Eq. (2.24). The summation is a geometric series.

2.2 Coherent States

We have shown that a phase state $|\phi\rangle$ has a well-defined phase. However, as a classical field, not only the phase but also the field amplitude should be well-defined, that is, we expect that $\langle \mathbf{E} \rangle$ does not vanish, and $\sigma(\mathbf{E})$ is small. Since the phase states are not normalized nor physical, we have to find other states.

The goal is to find the states $|\alpha\rangle$ such that the expectation of the electric field $\langle \alpha | \mathbf{E} | \alpha \rangle$ is proportional to the classical field $\mathcal{E}_\omega(\mathbf{r}) + \mathcal{E}_\omega^*(\mathbf{r})$. By observing that

$$\mathbf{E}_\omega(\mathbf{r}) = \frac{[\mathcal{E}_\omega(\mathbf{r})a + \mathcal{E}_\omega^*(\mathbf{r})a^\dagger]}{2}, \quad (2.29)$$

one finds that if the states $|\alpha\rangle$ are the eigenstates of the annihilation operator a ,

$$a|\alpha\rangle = \alpha|\alpha\rangle, \quad (2.30)$$

with the eigenvalues α , the expectation value $\langle \alpha | \mathbf{E} | \alpha \rangle$ is the same as the classical field. Since the operator a is not hermitian, the eigenvalues α can be complex numbers in general. It turns out that the states $|\alpha\rangle$, called “coherent states”, are the most classical states. Let’s find out the coherent states in the number state bases. We expand the coherent states as

$$|\alpha\rangle = \sum_n C_n |n\rangle, \quad (2.31)$$

and plug it in Eq. (2.30),

$$a|\alpha\rangle = \sum_n C_n a|n\rangle = \alpha \sum_n C_n |n\rangle \quad (2.32)$$

$$\Rightarrow \sum_n C_n \sqrt{n} |n-1\rangle = \alpha \sum_n C_n |n\rangle. \quad (2.33)$$

We obtain

$$C_{n+1} = \alpha \frac{C_n}{\sqrt{n+1}}, \quad (2.34)$$

$$C_n = \frac{\alpha^n}{\sqrt{n!}} C_0, \quad (2.35)$$

and thus

$$|\alpha\rangle = C_0 \sum_n \frac{\alpha^n}{\sqrt{n!}} |n\rangle. \quad (2.36)$$

The coefficient C_0 is fixed by the normalization condition,

$$\langle \alpha | \alpha \rangle = |C_0|^2 \sum_{m,n} \frac{\alpha^n (\alpha^*)^m}{\sqrt{n!m!}} \langle m | n \rangle, \quad (2.37)$$

where one finds

$$C_0 = e^{-\frac{|\alpha|^2}{2}}. \quad (2.38)$$

The coherent states are

$$|\alpha\rangle = e^{-\frac{|\alpha|^2}{2}} \sum_n \frac{\alpha^n}{\sqrt{n!}} |n\rangle \quad (2.39)$$

$$= e^{-\frac{|\alpha|^2}{2}} \sum_n \frac{\alpha^n (a^\dagger)^n}{n!} |0\rangle \quad (2.40)$$

$$= e^{-\frac{|\alpha|^2}{2}} e^{\alpha a^\dagger} |0\rangle. \quad (2.41)$$

Exercise 2: Normalization Constant

Show Eq. (2.38). Begin with Eq. (2.37).

The expectations are

$$\langle \alpha | \mathbf{E} | \alpha \rangle = \left\langle \alpha \left| \frac{[\mathcal{E}_\omega(\mathbf{r})a + \mathcal{E}_\omega^*(\mathbf{r})a^\dagger]}{2} \right| \alpha \right\rangle \quad (2.42)$$

$$= \text{Re}[\alpha \mathcal{E}_\omega(\mathbf{r})] \quad (2.43)$$

$$\begin{aligned} \langle \alpha | \text{abs}[\mathbf{E}]^2 | \alpha \rangle &= \left\langle \alpha \left| \frac{\text{abs}[\mathcal{E}_\omega(\mathbf{r})a + \mathcal{E}_\omega^*(\mathbf{r})a^\dagger]^2}{4} \right| \alpha \right\rangle \\ &= \text{abs}[\text{Re}[\alpha \mathcal{E}_\omega(\mathbf{r})]]^2 + \frac{|\mathcal{E}_\omega(\mathbf{r})|^2}{4}. \end{aligned} \quad (2.44)$$

The standard deviation of the electric field is

$$\sigma(\mathbf{E}) = \frac{|\mathcal{E}_\omega(\mathbf{r})|}{2}. \quad (2.45)$$

The standard deviation is relatively small compared to the field amplitude when $|\alpha|$ is large. We can see this by dividing $\sigma(\mathbf{E})$ with $\langle \alpha | \mathbf{E} | \alpha \rangle$,

$$\frac{\sigma(\mathbf{E})}{\langle \alpha | \mathbf{E} | \alpha \rangle} = \frac{|\mathcal{E}_\omega(\mathbf{r})|}{2\text{Re}[\alpha \mathcal{E}_\omega(\mathbf{r})]}. \quad (2.46)$$

The coherent states $|\alpha\rangle$ indeed have the minimum uncertainty. Using the quadrature operators X and Y , one can show that the coherent states have

$$\sigma(X) = \sigma(Y) = \frac{1}{2}. \quad (2.47)$$

Exercise 3: Uncertainty Relations

Show Eq. (2.47). Hints:

$$(a) \langle \alpha | X | \alpha \rangle = \frac{\alpha + \alpha^*}{2}$$

$$(b) \langle \alpha | X^2 | \alpha \rangle = \left(\frac{\alpha + \alpha^*}{2} \right)^2 + \frac{1}{4}. \text{ Note that } (a + a^\dagger)^2 = a^2 + 2a^\dagger a + (a^\dagger)^2 + 1$$

The physical meaning of α is the dimensionless amplitude, which is seen from that the average number \bar{n} of a coherent state $|\alpha\rangle$ is

$$\bar{n} = \langle \alpha | N | \alpha \rangle = \langle \alpha | a^\dagger a | \alpha \rangle = |\alpha|^2. \quad (2.48)$$

The standard deviation $\sigma(N)$ is

$$\sigma(N) = |\alpha| = \bar{n}^{\frac{1}{2}}. \quad (2.49)$$

The standard deviation $\sigma(N)$ over the average number \bar{n} is

$$\frac{\sigma(N)}{\bar{n}} = \bar{n}^{-\frac{1}{2}}. \quad (2.50)$$

The probability p_n of measuring the number state $|n\rangle$ is a Poisson distribution

$$p_n = |C_n|^2 = e^{-|\alpha|^2} \frac{|\alpha|^{2n}}{n!} = e^{-\bar{n}} \frac{\bar{n}^n}{n!}. \quad (2.51)$$

The phase distribution function $\mathcal{P}(\phi)$ of a coherent state is

$$\mathcal{P}(\phi) = \frac{e^{-|\alpha|^2}}{2\pi} \left| \sum_n \frac{\alpha^n}{\sqrt{n!}} \right|^2. \quad (2.52)$$

Let $\alpha = |\alpha| e^{i\phi}$. One can show that as $\bar{n} = |\alpha|^2$ is large, the distributions become approximately the Gaussian distributions (See Ref. [1]),

$$p_n \simeq (2\pi\bar{n})^{-1/2} e^{-\frac{(n-\bar{n})^2}{2\bar{n}}}, \quad (2.53)$$

$$\mathcal{P}(\phi) \simeq \sqrt{\frac{2\bar{n}}{\pi}} e^{-2\bar{n}(\phi-\bar{\phi})^2}. \quad (2.54)$$

2.3 Displaced Vacuum States

The physical meaning of α is the dimensionless (complex) amplitude of a coherent state. The vacuum state is indeed a coherent state in the limit $\alpha \rightarrow 0$. Conversely, a coherent state is obtained by changing the complex amplitude α of the vacuum state. Mathematically, such a shift of α is done by the displacement operator $D(\alpha)$,

$$|\alpha\rangle = D(\alpha)|0\rangle. \quad (2.55)$$

The displacement operator $D(\alpha)$ has the explicit form

$$D(\alpha) = \exp(\alpha a^\dagger - \alpha^* a). \quad (2.56)$$

To show this, first consider the special case of Baker–Campbell–Hausdorff formula, if

$$[A, [A, B]] = [B, [A, B]] = 0, \quad (2.57)$$

we have

$$e^{A+B} = e^{-\frac{1}{2}[A,B]} e^A e^B \quad (2.58)$$

$$= e^{\frac{1}{2}[B,A]} e^B e^A. \quad (2.59)$$

With $A = \alpha a^\dagger$, $B = -\alpha^* a$, and $[A, B] = |\alpha|^2$, the displacement operator $D(\alpha)$ becomes

$$D(\alpha) = e^{-\frac{|\alpha|^2}{2}} e^{\alpha a^\dagger} e^{-\alpha^* a}. \quad (2.60)$$

Using the relations

$$e^{-\alpha^* a} |0\rangle = \left(\mathbb{1} - \alpha^* a + \frac{(-\alpha^* a)^2}{2!} + \dots \right) |0\rangle = |0\rangle, \quad (2.61)$$

we obtain

$$D(\alpha) |0\rangle = e^{-\frac{|\alpha|^2}{2}} e^{\alpha a^\dagger} e^{-\alpha^* a} |0\rangle \quad (2.62)$$

$$= e^{-\frac{|\alpha|^2}{2}} e^{\alpha a^\dagger} |0\rangle \quad (2.63)$$

$$= e^{-\frac{|\alpha|^2}{2}} e^{\alpha a^\dagger} |0\rangle \quad (2.64)$$

$$= e^{-\frac{|\alpha|^2}{2}} \sum_n \frac{\alpha^n (a^\dagger)^n}{n!} |0\rangle \quad (2.65)$$

$$= e^{-\frac{|\alpha|^2}{2}} \sum_n \frac{\alpha^n}{\sqrt{n!}} |n\rangle \quad (2.66)$$

$$= |\alpha\rangle. \quad (2.67)$$

The displacement operator $D(\alpha)$ is unitary and satisfies the relation

$$D(\alpha) D^\dagger(\alpha) = D^\dagger(\alpha) D(\alpha) = \mathbb{1}, \quad (2.68)$$

$$D^\dagger(\alpha) = D(-\alpha). \quad (2.69)$$

The displacement operators satisfy the law of addition; operations by two subsequent displacement operator $D(\alpha)$ and $D(\beta)$ give a total displacement operator

$$D(\alpha) D(\beta) = e^{i\text{Im}[\alpha\beta^*]} D(\alpha + \beta). \quad (2.70)$$

We see that the total displacement is $\alpha + \beta$, that is, the sum of the displacements of the individual displacement operators. An extra phase $\text{Im}[\alpha\beta^*]$ is the quantum feature, and note that although the total displacement does not depend on the order of the operators, the phase does depend.

Note 3: Displacement Operator

For now, a displacement operator is just a mathematical tool. Later, as we learn light-matter interaction, we will know that a displacement operator is the evolution operator of a sinusoidal driving source, $\mathcal{H}_i(t) \sim \sin(\omega t + \phi)$. That is, if we turn on a sinusoidal driving source, the vacuum state will be shifted in the complex α space. This is one method to generate coherent

states.

2.4 Dynamics of Coherent States

The dynamics of a coherent state $|\alpha\rangle$ is given by the Schrödinger's picture,

$$|\alpha(t)\rangle = e^{-i\frac{\hat{H}t}{\hbar}}|\alpha(0)\rangle \quad (2.71)$$

$$= e^{-i\frac{\omega}{2}t} e^{-\frac{|\alpha|^2}{2}} \sum_n \frac{\alpha^n e^{-in\omega t}}{\sqrt{n!}} |n\rangle \quad (2.72)$$

$$= e^{-i\frac{\omega}{2}t} |\alpha(0)\rangle e^{-i\omega t}. \quad (2.73)$$

Thus, the amplitude $\alpha(t)$ is

$$\alpha(t) = \alpha(0)e^{-i\omega t}. \quad (2.74)$$

Although every photon mode $\mathcal{E}_\omega(\mathbf{r})$ can be quite different from one system to another system, we can use the dimensionless quadrature operators \hat{X} and \hat{Y} to describe the dynamics. Recall that \hat{X} is analogous to the position operator, and \hat{Y} is analogous to the momentum operator. We can express a coherent state in the X basis,

$$\psi_\alpha(X) = \langle X|\alpha\rangle, \quad (2.75)$$

where $|X\rangle$ is the eigenvector of X

$$\hat{X}|X\rangle = X|X\rangle. \quad (2.76)$$

To find $\psi_\alpha(X)$, we begin with

$$\langle X|\hat{a}|\alpha\rangle = \alpha\langle X|\alpha\rangle \quad (2.77)$$

$$\Rightarrow \langle X|\hat{X} + i\hat{Y}|\alpha\rangle = \alpha\langle X|\alpha\rangle \quad (2.78)$$

$$\Rightarrow \left(X + \frac{\partial}{\partial X}\right)\langle X|\alpha\rangle = \alpha\langle X|\alpha\rangle \quad (2.79)$$

$$\Rightarrow \frac{\partial\psi_\alpha(X)}{\partial X} = (\alpha - X)\psi_\alpha(X) \quad (2.80)$$

$$\Rightarrow \psi_\alpha(X) = \sqrt{\frac{2}{\pi}} e^{-\frac{(X - \text{Re}[\alpha])^2}{2}} e^{i\text{Im}[\alpha]X}, \quad (2.81)$$

where we have used the normalization condition to derive the last step. The wavefunction $\psi_\alpha(X)$ is a Gaussian distribution, and its peak position is

$$X_p(t) = \text{Re}[\alpha(t)] \quad (2.82)$$

$$= |\alpha(0)| \cos(\phi_0 - \omega t). \quad (2.83)$$

with $\alpha(0) = |\alpha(0)|e^{i\phi_0}$. The peak position $X_p(t)$ is the same as that of a classical harmonic oscillator. Also the wavefunction $\psi_\alpha(X)$ has a minimum spread of X and P . Thus, a coherent state is the most classical state.

Summary 1: Coherent States

Coherent states are

- eigenstates of the annihilation operator a .
- displaced vacuum states.
- most classical states whose phase and amplitude distributions are narrow.
- most classical states whose X and Y distributions are narrow.
- minimum uncertainty states.

2.5 Properties of Coherent States

2.5.1 Orthogonality

Two coherent states $|\alpha\rangle$ and $|\beta\rangle$ are not orthogonal,

$$\langle\beta|\alpha\rangle = e^{-\frac{(|\alpha|^2+|\beta|^2)}{2}} \sum_{n,m} \frac{(\beta^*)^m (\alpha)^n}{\sqrt{m!n!}} \langle m|n\rangle \quad (2.84)$$

$$= e^{-\frac{(|\alpha|^2+|\beta|^2)}{2}} \sum_n \frac{(\beta^*)^n (\alpha)^n}{n!} \quad (2.85)$$

$$= e^{-\frac{(|\alpha|^2+|\beta|^2)}{2}} e^{\beta^* \alpha} \quad (2.86)$$

$$= e^{-\frac{|\alpha-\beta|^2}{2}} e^{\frac{\beta^* \alpha - \beta \alpha^*}{2}}, \quad (2.87)$$

which does not vanish.

2.5.2 Identity

The identity can be expressed with the coherent states,

$$\int \frac{d^2\alpha}{\pi} |\alpha\rangle\langle\alpha| \equiv \int \frac{d\text{Re}[\alpha]d\text{Im}[\alpha]}{\pi} |\alpha\rangle\langle\alpha| = \mathbb{1}. \quad (2.88)$$

Derivation 2: Identity with Coherent States

The proof of Eq. (2.88) is as follows. Let $\alpha = re^{i\phi}$ and.

$$d\alpha^2 = d\text{Re}[\alpha]d\text{Im}[\alpha] = r dr d\phi. \quad (2.89)$$

The left hand side of Eq. (2.88) becomes

$$\int \frac{rdrd\phi}{\pi} |\alpha\rangle\langle\alpha| = \int \frac{rdrd\phi}{\pi} e^{-r^2} \sum_{m,n} \frac{e^{i(n-m)\phi} r^{m+n}}{n!} |m\rangle\langle n| \quad (2.90)$$

$$= \sum_n \frac{\int dr e^{-r^2} 2r^{2n+1}}{n!} |n\rangle\langle n| \quad (2.91)$$

$$= \sum_n \frac{\int du e^{-u} u^n}{n!} |n\rangle\langle n| \quad (2.92)$$

$$= \sum_n |n\rangle\langle n| = \mathbb{1}. \quad (2.93)$$

2.5.3 Coherent State Representations of Operators

Any operator X can be expressed in the coherent state bases with the identity Eq. (2.88),

$$X = \int \frac{d^2\alpha}{\pi} \int \frac{d^2\beta}{\pi} |\alpha\rangle\langle\alpha| X |\beta\rangle\langle\beta|. \quad (2.94)$$

However, coherent states are not orthogonal, so the coherent states form an **over-complete** set of bases.¹ It is possible to write X in the coherent state diagonal form.

An operator X is uniquely determined by $\langle\alpha|X|\alpha\rangle$. The diagonal element $\langle\alpha|X|\alpha\rangle$ in the number state basis is

$$\langle\alpha|X|\alpha\rangle = \exp(-|\alpha|^2) \sum_{m,n} \frac{\langle n|X|m\rangle \alpha^m (\alpha^*)^n}{\sqrt{m!n!}}, \quad (2.95)$$

indicating that $\langle\alpha|X|\alpha\rangle$ contains all the information of the elements $\langle n|X|m\rangle$, which forms a complete set.

Coherent state diagonal representation. Suppose that X has a series expansion of a and a^\dagger in the antinormal ordering,

$$X = \sum_{mn} \chi_{nm}^A a^n (a^\dagger)^m, \quad (2.96)$$

where χ_{nm}^A is a c -number. The subscript A denotes the antinormal ordering. Inserting the identity, we obtain

$$X = \sum_{mn} \chi_{nm}^A a^n \left(\int \frac{d^2\alpha}{\pi} |\alpha\rangle\langle\alpha| \right) (a^\dagger)^m \quad (2.97)$$

$$= \int d^2\alpha \chi^A(\alpha) |\alpha\rangle\langle\alpha| \quad (2.98)$$

¹See Se. 5.4. of Ref. [2] for a more rigorous discussion.

where

$$\chi^A(\alpha) = \frac{1}{\pi} \sum_{mn} \chi_{nm}^A \alpha^n (\alpha^*)^m, \quad (2.99)$$

is a c -number.

3 Phase Space Distributions

Given a density matrix ρ , there are three important distribution functions which are the quantum analogs of the classical probability density $f(x, p)$.

3.1 Wigner Distribution

The Wigner function $W(\alpha)$ is defined as

$$W(\alpha) = \int \frac{d^2\eta}{\pi^2} e^{\eta^* \alpha - \eta \alpha^*} \chi_W(\eta), \quad (3.1)$$

where the characteristic function $\chi_W(\eta)$ is

$$\chi_W(\eta) = \text{Tr} \left[\rho e^{\eta a^\dagger - \eta^* a} \right]. \quad (3.2)$$

Exercise 4: Normalization

Show that

$$\int \frac{d^2\alpha}{\pi^2} e^{\eta^* \alpha - \eta \alpha^*} = \delta_2(\eta) \equiv \delta(\text{Re}[\eta]) \delta(\text{Im}[\eta]), \quad (3.3)$$

and use the result and Eq. (3.1) to show

$$\int d^2\alpha W(\alpha) = 1. \quad (3.4)$$

Hint: a delta function can be expressed as

$$\delta(x) = \frac{1}{2\pi} \int e^{ikx} dk. \quad (3.5)$$

Hint: let $\alpha = x + iy$ and use the identity $\delta(x) = \frac{1}{2\pi} \int e^{iqx} dq$.

The ensemble average of an operator X in this representation is

$$\langle X \rangle = \int d^2\alpha \chi^W(\alpha) W(\alpha), \quad (3.6)$$

where

$$\chi^W(\alpha) = \sum_{n,m} \chi_{nm}^W \alpha^n (\alpha^*)^m \quad (3.7)$$

The coefficient χ_{nm}^W is the Weyl(symmetric)-ordering representation of an operator X ,

$$X = \sum_{m,n} \chi_{nm}^W \left(\frac{(a^\dagger)^n a^m + a^m (a^\dagger)^n}{2} \right). \quad (3.8)$$

3.2 Glauber–Sudarshan P -function

The P -function is defined by

$$\rho = \int d^2\alpha P(\alpha) |\alpha\rangle\langle\alpha|, \quad (3.9)$$

and satisfies the normalization condition

$$1 = \text{Tr}[\rho] = \int d^2\alpha P(\alpha). \quad (3.10)$$

The P -function can be obtained by the normal-ordering characteristic function

$$P(\alpha) = \int \frac{d^2\eta}{\pi^2} e^{\eta^* \alpha - \eta \alpha^*} \chi_N(\eta), \quad (3.11)$$

where the characteristic function $\chi_N(\eta)$ is

$$\chi_N(\eta) = \text{Tr} \left[\rho e^{\eta a^\dagger} e^{-\eta^* a} \right]. \quad (3.12)$$

The ensemble average of an operator X in this representation is

$$\langle X \rangle = \int d^2\alpha \chi^N(\alpha) P(\alpha), \quad (3.13)$$

where

$$\chi^N(\alpha) = \sum_{n,m} \chi_{nm}^N (\alpha^*)^n \alpha^m \quad (3.14)$$

The coefficient χ_{nm}^N is the normal-ordering representation of an operator X ,

$$X = \sum_{m,n} \chi_{nm}^N (a^\dagger)^n a^m. \quad (3.15)$$

Note 4: Classical and Nonclassical States

A state with $P(\alpha) < 0$ is defined as a nonclassical state.

3.3 Q-function

The Q-function is defined by

$$Q(\alpha) = \frac{1}{\pi} \langle \alpha | \rho | \alpha \rangle, \quad (3.16)$$

which is always positive since it is the diagonal element of the density matrix. The function $Q(\alpha)$ satisfies

$$0 \leq Q(\alpha) \leq \frac{1}{\pi}, \quad (3.17)$$

and

$$\text{Tr}[\rho] = \int d^2\alpha Q(\alpha) = 1. \quad (3.18)$$

The Q-function can be obtained by the antinormal-ordering characteristic function

$$Q(\alpha) = \int \frac{d^2\eta}{\pi^2} e^{\eta^* \alpha - \eta \alpha^*} \chi_A(\eta), \quad (3.19)$$

where the characteristic function $\chi_A(\eta)$ is

$$\chi_A(\eta) = \text{Tr} \left[\rho e^{-\eta^* a} e^{\eta a^\dagger} \right]. \quad (3.20)$$

The ensemble average of an operator X in this representation is

$$\langle X \rangle = \int d^2\alpha \chi^A(\alpha) Q(\alpha), \quad (3.21)$$

where

$$\chi^A(\alpha) = \sum_{n,m} \chi_{nm}^A \alpha^n (\alpha^*)^m. \quad (3.22)$$

The coefficient χ_{nm}^A is the antinormal-ordering representation of an operator X ,

$$X = \sum_{m,n} \chi_{nm}^A a^n (a^\dagger)^m. \quad (3.23)$$

	$W(\alpha)$	$Q(\alpha)$	$P(\alpha)$
coherent state $ \alpha_0\rangle$	$\frac{2}{\pi} e^{-2 \alpha-\alpha_0 ^2}$	$\frac{1}{\pi} e^{- \alpha-\alpha_0 ^2}$	$\delta^2(\alpha - \alpha_0)$
thermal ensemble	$\frac{1}{\pi(\bar{n}+1/2)} \exp\left(-\frac{ \alpha ^2}{\bar{n}+1/2}\right)$	$\frac{1}{\pi(\bar{n}+1)} \exp\left(-\frac{ \alpha ^2}{\bar{n}+1}\right)$	$\frac{1}{\pi(\bar{n})} \exp\left(-\frac{ \alpha ^2}{\bar{n}}\right)$
pure ensemble $ 1\rangle\langle 1 $	$-(1-4 \alpha ^2)\frac{2}{\pi} e^{-2 \alpha ^2}$	$\frac{ \alpha ^2}{\pi} e^{- \alpha ^2}$	singular

Table 1: Examples of $W(\alpha)$, $Q(\alpha)$, and $P(\alpha)$

Summary 2: Coherent States

- (a) The phase space of photon states or ensembles are described by the two dimensional complex α plane.
- (b) The real part and imaginary part of α are related to the quadrature operator X and Y .

$$X = \text{Re}[\alpha], \quad (3.24)$$

$$Y = \text{Im}[\alpha]. \quad (3.25)$$

- (c) A coherent state $|\alpha_0\rangle$ is a fuzzy circle on the complex α plane.
- (d) The coherent states are not orthogonal, so they are overcomplete.
- (e) There are three ways to write the probability density
- Wigner distribution $W(\alpha)$: symmetric ordering
 - Q -function $Q(\alpha)$: antinormal ordering
 - P -function $P(\alpha)$: normal ordering

References

- [1] S. M. Barnett and D. T. Pegg, *J. Mod. Opt.*, 36 (1989), 7.
- [2] J. C. Garrison and R. Y. Chiao, *Quantum Optics*, Oxford University Press 2008